

KLEINIAN GROUPS WHICH ARE ALMOST FUCHSIAN

FRANCIS BONAHO

ABSTRACT. We consider the space of all quasifuchsian metrics on the product of a surface with the real line. We show that, in a neighborhood of the submanifold consisting of fuchsian metrics, every non-fuchsian metric is completely determined by the bending data of its convex core.

Let S be a surface of finite topological type, obtained by removing finitely many points from a compact surface without boundary, and with negative Euler characteristic. We consider complete hyperbolic metrics on the product $S \times]-\infty, \infty[$.

The simplest ones are the fuchsian metrics defined as follows. Because of our hypothesis that the Euler characteristic of S is negative, S admits a finite area hyperbolic metric, for which S is isometric to the quotient of the hyperbolic plane $\mathbb{H}^2$ by a discrete group Γ of isometries. The group Γ uniquely extends to a group of isometries of the hyperbolic 3-space $\mathbb{H}^3$ respecting the transverse orientation of $\mathbb{H}^2 \subset \mathbb{H}^3$, for which the quotient $\mathbb{H}^3/\Gamma$ has a natural identification with $S \times]-\infty, \infty[$. A *fuchsian metric* is any metric on $S \times]-\infty, \infty[$ obtained in this way. Note that the image of $\mathbb{H}^2$ in $\mathbb{H}^3$ provides in this case a totally geodesic surface in $S \times]-\infty, \infty[$, isometric to the original metric on S .

These examples can be perturbed to more complex hyperbolic metrics on $S \times]-\infty, \infty[$. See for instance [Th1][Mas]. A *quasifuchsian metric* on $S \times]-\infty, \infty[$ is one which is obtained by quasi-conformal deformation of a fuchsian metric. Equivalently, a quasifuchsian metric is a geometrically finite hyperbolic metric on $S \times]-\infty, \infty[$ whose cusps exactly correspond to the ends of S . These also correspond to the interior points in the space of all hyperbolic metrics on $S \times]-\infty, \infty[$ for which the ends of S are parabolic [Mar][Su].

If m is a quasifuchsian metric on $S \times]-\infty, \infty[$, the totally geodesic copy of S which occurred in the fuchsian case is replaced by the *convex core* $C(m)$, defined as the smallest non-empty closed m -convex subset of $S \times]-\infty, \infty[$. If m is not fuchsian, $C(m)$ is 3-dimensional and its boundary consists of two copies of S , each facing an end of $S \times]-\infty, \infty[$. The geometry of $\partial C(m)$ was investigated by Thurston [Th1]; see also [EpM]. The component of $\partial C(m)$ that faces the end $S \times \{+\infty\}$ is a pleated surface, totally geodesic almost everywhere, but bent along a family of simple geodesics; this bending is described and quantified by a measured geodesic lamination $\beta^+(m)$ on S . Similarly, the bending of the negative component of $\partial C(m)$, namely the one facing $S \times \{-\infty\}$, is determined by a measured geodesic lamination $\beta^-(m)$.

If $\mathcal{Q}(S)$ denotes the space of isotopy classes of quasifuchsian metrics on $S \times]-\infty, \infty[$ and if $\mathcal{ML}(S)$ is the space of measured geodesic laminations on S , the

Date: February 1, 2008.

This work was partially supported by grant DMS-0103511 from the National Science Foundation.

rule $m \mapsto (\beta^+(m), \beta^-(m))$ defines a map $\beta : \mathcal{Q}(S) \rightarrow \mathcal{ML}(S) \times \mathcal{ML}(S)$. By definition, $\beta(m) = (0, 0)$ if the metric m is fuchsian, in which case the convex core $C(m)$ is just a totally m -geodesic copy of S . The image $\beta(m) \in \mathcal{ML}(S)^2$, interpreted as a measured geodesic lamination on two copies of S , is the *bending measured geodesic lamination* of the quasifuchsian metric m .

The space $\mathcal{Q}(S)$ is a manifold of dimension 2θ where, if $\chi(S)$ is the Euler characteristic of S and p is its number of ends, $\theta = -3\chi(S) - p \geq 0$. It contains the space $\mathcal{F}(S)$ as a proper submanifold of dimension θ . To some extent, the bending measured lamination $\beta(m)$ measures how far the metric $m \in \mathcal{Q}(S)$ is from being fuchsian. Finally, recall that the space $\mathcal{ML}(S)$ of measured geodesic laminations is a piecewise linear manifold of dimension θ .

Thurston conjectured that the restriction of the bending map to β could be used to parametrize $\mathcal{Q}(S) - \mathcal{F}(S)$, namely that it induces a homeomorphism between $\mathcal{Q}(S) - \mathcal{F}(S)$ and an open subset of $\mathcal{ML}(S)^2$. The image of β was determined in [BoO].

The goal of the current paper is to prove Thurston's conjecture on a neighborhood of the space of Fuchsian metrics.

Theorem 1. *There exists an open neighborhood V of the fuchsian submanifold $\mathcal{F}(S)$ in $\mathcal{Q}(S)$ such that the bending map $\beta : \mathcal{Q}(S) \rightarrow \mathcal{ML}(S)^2$ induces a homeomorphism between $V - \mathcal{F}(S)$ and its image.*

There are well-known restrictions for $(\mu, \nu) \in \mathcal{ML}(S)^2$ to be in the image of β ; see for instance [BoO]. In particular, if $(\mu, \nu) \neq (0, 0)$ is the bending measured lamination of some $m \in \mathcal{Q}(S)$, then the measured geodesic laminations μ and ν must *fill up* the surface S , in the sense that every non-trivial measured geodesic lamination has non-zero geometric intersection number with at least one of μ, ν . This is equivalent to the condition that every component of $S - \mu \cup \nu$ is, either a topological disk bounded by the union of finitely many geodesic arcs, or a topological annulus bounded on one side by the union of finitely many geodesic arcs and going to a cusp on the other side.

Let $\mathcal{FML}(S)$ denote the open subset of $\mathcal{ML}(S)^2$ consisting of those (μ, ν) where μ and ν fill up S . Note that $\mathcal{FML}(S)$ is endowed with an action of $\mathbb{R}^+$, defined by $t(\mu, \nu) = (t\mu, t\nu)$, which decomposes $\mathcal{FML}(S)$ as the union of pairwise disjoint rays (=orbits) $]0, \infty[(\mu, \nu)$.

Theorem 2. *In Theorem 1, the neighborhood V of $\mathcal{F}(S)$ and its image $U = \beta(V)$ can be chosen so that $U - \{(0, 0)\}$ is an open subset of $\mathcal{FML}(S)$ which intersects each ray $]0, \infty[(\mu, \nu)$ in an interval $]0, \varepsilon_{\mu\nu}[(\mu, \nu)$.*

Theorems 1 and 2 are proved later as Theorem 14. The main idea of the proof is to construct an inverse $\beta^{-1} : U - \{(0, 0)\} \rightarrow V - \mathcal{F}(S)$, and splits into two steps: an infinitesimal part, and a transversality argument based on the infinitesimal part. The infinitesimal part is now relatively classical; see for instance [Se1]. There are restrictions on which bending data can be realized by an infinitesimal deformation of $m_0 \in \mathcal{F}(S)$. Through the complex structure of $\mathcal{Q}(S)$, where multiplication by $i = \sqrt{-1}$ converts bending to shearing, these restrictions can be expressed in a purely 2-dimensional context; see Section 2. The main part of the proof is to show by a transversality argument that any infinitesimal bending data can actually be realized by a deformation. The only significant idea of the paper is to apply the transversality argument, not in $\mathcal{Q}(S)$ where the necessary hypotheses are not

realized, but in the manifold-with-boundary $\check{\mathcal{Q}}(S)$ obtained by blowing up $\mathcal{Q}(S)$ along the fuchsian submanifold $\mathcal{F}(S)$.

Switching to the blow-up manifold $\check{\mathcal{Q}}(S)$ actually provides a better understanding of the restriction of β on a neighborhood of $\mathcal{F}(S)$ and of its inverse. See Theorem 13 for a precise statement.

Acknowledgements: There is strong evidence that the content of this paper has been known to Bill Thurston for about twenty years. I personally learned much of the material of Section 2 many years ago from David Epstein who, I believe, had learned it from Thurston. I am also grateful to Pete Storm and Dick Canary for helping me to clarify my thoughts.

1. THE EARTHQUAKE SECTION

We consider the *Teichmüller space* $\mathcal{T}(S)$, namely the space of isotopy classes of finite area complete hyperbolic metrics on the surface S . Recall that $\mathcal{T}(S)$ is diffeomorphic to $\mathbb{R}^\theta$, where $\theta = -3\chi(S) - p \geq 0$ if $\chi(S)$ is the Euler characteristic of S and p is its number of ends.

A standard deformation of a metric $m \in \mathcal{T}(S)$ is the *left earthquake* $E_m^\mu \in \mathcal{T}(S)$ along the measured geodesic lamination μ , as constructed in [Th2][Ke1][EpM]. We consider the *infinitesimal left earthquake vector* $e_m^\mu = \frac{d}{dt}E_m^{t\mu}|_{t=0} \in T_m\mathcal{T}(S)$. This provides a section $e^\mu : \mathcal{T}(S) \rightarrow T\mathcal{T}(S)$ of the tangent bundle of $\mathcal{T}(S)$, defined by $m \mapsto e_m^\mu$.

There similarly exists a *right earthquake* $E_m^{-\mu}$ along μ ; the notation is justified by the fact that $m \mapsto E_m^{-\mu}$ is the inverse of $m \mapsto E_m^\mu$. We can then consider the *infinitesimal right earthquake vector* $e_m^{-\mu} = \frac{d}{dt}E_m^{-t\mu}|_{t=0} \in T_m\mathcal{T}(S)$ and the corresponding section $e^{-\mu} : \mathcal{T}(S) \rightarrow T\mathcal{T}(S)$ of the tangent bundle of $\mathcal{T}(S)$. Note that $e_m^{-\mu} = -e_m^\mu$, but it will be convenient to keep a separate notation.

Recall that the measured geodesic laminations $\mu, \nu \in \mathcal{ML}(S)$ *fill up* the surface S if every non-trivial measured geodesic lamination has non-zero geometric intersection number with at least one of μ, ν . This is equivalent to the condition that every component of $S - \mu \cup \nu$ is, either a topological disk bounded by the union of finitely many geodesic arcs, or a topological annulus bounded on one side by the union of finitely many geodesic arcs and going to a cusp on the other side.

Proposition 3. *Let $\mu, \nu \in \mathcal{ML}(S)$ be two non-zero measured geodesic laminations. The intersection of the two sections $e^\mu, e^{-\nu} : \mathcal{T}(S) \rightarrow T\mathcal{T}(S)$ of the tangent bundle of $\mathcal{T}(S)$ is transverse. These sections meet in exactly one point if μ and ν fill up the surface S , and are otherwise disjoint.*

Proof. We first translate the problem in terms of the length functions $l_\mu, l_\nu : \mathcal{T}(S) \rightarrow \mathbb{R}$ which to a metric $m \in \mathcal{T}(S)$ associate the m -lengths of the measured geodesic lamination μ and ν . The Weil-Petersson symplectic form on $\mathcal{T}(S)$ induces an isomorphism between its tangent bundle $T\mathcal{T}(S)$ and its cotangent bundle $T^*\mathcal{T}(S)$. A celebrated result of Scott Wolpert [Wol] asserts that this isomorphism sends the section e^μ of $T\mathcal{T}(S)$ to the section dl_μ of $T^*\mathcal{T}(S)$. Therefore, Proposition 3 is equivalent to showing that the sections dl_μ and $-dl_\nu$ transversely meet in 1 or 0 point, according to whether μ and ν fill up the surface S or not.

First consider the case where μ and ν fill up S . The intersection of the section dl_μ and $-dl_\nu$ of $T^*\mathcal{T}(S)$ correspond to the points $m \in \mathcal{T}(S)$ where $d_m l_\mu = -d_m l_\nu$, namely to the critical points of the function $l_\mu + l_\nu : \mathcal{T}(S) \rightarrow \mathbb{R}$. It is proved in

[Ke1][Ke2] that, because μ and ν fill up S , the function $l_\mu + l_\nu$ admits a unique critical point m_0 . In addition, the hessian of $l_\mu + l_\nu$ at m_0 is positive definite [Wo2][Ke2].

Let $u = d_{m_0}l_\mu = -d_{m_0}l_\nu \in T_{m_0}^*\mathcal{T}(S)$. The intersection of the tangent spaces of the sections dl_μ and $-dl_\nu$ at u consists of all the vectors of the form $T_{m_0}(dl_\mu)(v) = T_{m_0}(-dl_\nu)(v)$ for some $v \in T_{m_0}\mathcal{T}(S)$, where $T_{m_0}(dl_\mu), T_{m_0}(-dl_\nu) : T_{m_0}\mathcal{T}(S) \rightarrow T_u T^*\mathcal{T}(S)$ denote the tangent maps of the sections $dl_\mu, -dl_\nu : \mathcal{T}(S) \rightarrow T^*\mathcal{T}(S)$.

Note that, for an arbitrary $w \in T_{m_0}\mathcal{T}(S)$, the vectors $T_{m_0}(dl_\mu)(w)$ and $T_{m_0}(-dl_\nu)(w)$ both project to w through the tangent map of the projection $T^*\mathcal{T}(S) \rightarrow \mathcal{T}(S)$. In particular, the difference $T_{m_0}(dl_\mu)(w) - T_{m_0}(-dl_\nu)(w)$ is tangent to the fiber, namely is an element of $T_{m_0}^*\mathcal{T}(S) \subset T_u T^*\mathcal{T}(S)$. Using local coordinates, one easily sees that this difference is the image of w under the homomorphism $T_{m_0}\mathcal{T}(S) \rightarrow T_{m_0}^*\mathcal{T}(S)$ induced by the hessian of the function $l_\mu + l_\nu$. (Beware of optimistic simplifications, though: for instance, $T_{m_0}(-dl_\nu)$ is not the same as $-T_{m_0}(dl_\nu)$ since they take their values in different tangent spaces.) Since the hessian is non-degenerate, this homomorphism is actually an isomorphism.

A consequence of this analysis is that, if $T_{m_0}(dl_\mu)(v) = T_{m_0}(-dl_\nu)(v)$, then v must be 0. Therefore, the intersection of the two sections dl_μ and $-dl_\nu$ of $T^*\mathcal{T}(S)$ at the point u is transverse, since these sections have dimension θ and the total space $T^*\mathcal{T}(S)$ has dimension 2θ . By Weil-Petersson duality, this proves that the sections e^μ of $e^{-\nu}$ of $T\mathcal{T}(S)$ have a transverse intersection consisting of exactly one point.

We now consider the case where μ and ν do not fill up the surface. In this case, the function $l_\mu + l_\nu$ has no critical point [Ke2] (see also the discussion in [Se1, §4.2]). By the above translation, it follows that the sections $e^\mu, e^{-\nu}$ are disjoint.

This concludes the proof of Proposition 3. $\square$

If the measured geodesic laminations μ, ν fill up the surface S , let $\kappa(\mu, \nu)$ denote the (unique) critical point of the length function $l_\mu + l_\nu : \mathcal{T}(S) \rightarrow \mathbb{R}$. As above, $\kappa(\mu, \nu)$ is also the unique $m \in \mathcal{T}(S)$ such that $e_m^\mu = e_m^{-\nu}$.

Lemma 4. *If $\kappa(\mu', \nu) = \kappa(\mu, \nu)$, then $\mu' = \mu$.*

Proof. By an (easy) infinitesimal version of [Th2] (see also [Ke2]), an infinitesimal left earthquake completely determines the measured geodesic lamination along which it is performed. If $e_m^{\mu'} = e_m^{-\nu} = e_m^\mu$, it follows that $\mu' = \mu$. $\square$

We will need a result similar to Proposition 3 in the unit tangent bundle $T^1\mathcal{T}(S)$. Recall that the fibers $T_m^1\mathcal{T}(S)$ of this bundle are the quotient of $T_m\mathcal{T}(S) - \{0\}$ under the equivalence relation which identifies v to tv when $t \in]0, \infty[$. In particular, $T^1\mathcal{T}(S)$ is a manifold of dimension $2\theta - 1$. Let $\bar{e}^\mu, \bar{e}^{-\nu} : \mathcal{T}(S) \rightarrow T^1\mathcal{T}(S)$ be the sections induced by e^μ and $e^{-\nu}$.

Proposition 5. *Let $\mu, \nu \in \mathcal{ML}(S)$ be two non-zero measured geodesic laminations. The intersection of the two sections $\bar{e}^\mu, \bar{e}^{-\nu} : \mathcal{T}(S) \rightarrow T^1\mathcal{T}(S)$ of the unit tangent bundle of $\mathcal{T}(S)$ is transverse. If μ and ν fill up the surface S , these sections meet along a section above a line $K(\mu, \nu)$ properly embedded in $\mathcal{T}(S)$. If μ, ν do not fill up S , the intersection is empty.*

Proof. The two sections meet above $m \in \mathcal{T}(S)$ if $\bar{e}_m^\mu = \bar{e}_m^{-\nu}$, namely if there is a $t > 0$ such that $e_m^{-\nu} = te_m^\mu = e_m^{t\mu}$. By Proposition 3, this can occur only when $t\mu$

and ν fill the surface, namely only when μ and ν fill the surface. Consequently, the two sections have empty intersection if μ and ν do not fill up the surface.

If μ and ν fill up the surface then, for every $t > 0$, Proposition 3 shows that there is a unique $m = \kappa(t\mu, \nu) \in \mathcal{T}(S)$ such that $e_m^{t\mu} = e_m^{-\nu}$. As a consequence, the two sections $\bar{e}^\mu, \bar{e}^{-\nu} : \mathcal{T}(S) \rightarrow T^1\mathcal{T}(S)$ meet exactly above the image $K(\mu, \nu)$ of the map $]0, \infty[\rightarrow \mathcal{T}(S)$ defined by $t \mapsto \kappa(t\mu, \nu)$.

If $m = \kappa(t\mu, \nu)$ so that $e_m^{t\mu} = e_m^{-\nu}$, the tangent space $T_{e_m^{t\mu}}T\mathcal{T}(S)$ is the sum of the tangent spaces $T_m e^{t\mu}(T_m\mathcal{T}(S))$ and $T_m e^{-\nu}(T_m\mathcal{T}(S))$ of the sections $e^{t\mu}$ and $e^{-\nu}$, by the transversality property of Proposition 3. Therefore, in the unit tangent bundle, the tangent space $T_{\bar{e}_m^\mu}T^1\mathcal{T}(S)$ is the sum of $T_m \bar{e}^\mu(T_m\mathcal{T}(S))$ and $T_m \bar{e}^{-\nu}(T_m\mathcal{T}(S))$. As a consequence, the intersection of the two sections $\bar{e}^\mu$ and $\bar{e}^{-\nu}$ is transverse above the point $m = \kappa(t\mu, \nu)$.

By transversality, the intersection of the sections is a submanifold of the image of $\bar{e}^\mu$. Its dimension is equal to 1, by consideration of the dimensions of $\mathcal{T}(S)$ and $T^1\mathcal{T}(S)$. Since the projection $\bar{e}^\mu(\mathcal{T}(S)) \rightarrow \mathcal{T}(S)$ is a diffeomorphism, it follows that the projection $K(\mu, \nu)$ of this projection is a 1-dimensional submanifold of $\mathcal{T}(S)$.

By definition, $\kappa(t\mu, \nu)$ is the unique minimum of the convex function $t dl_\mu + dl_\nu$, which has positive hessian at this minimum. It follows that $\kappa(t\mu, \nu)$ is a continuous function of t . Conversely, t is completely determined by $m = \kappa(t\mu, \nu)$ by Lemma 4. As a consequence, if $m = \kappa(t\mu, \nu)$ stays in a bounded subset of $\mathcal{T}(S)$, then t stays in a compact subset of $]0, \infty[$. In other words, the map $]0, \infty[\rightarrow \mathcal{T}(S)$ defined by $t \mapsto \kappa(t\mu, \nu)$ is injective, continuous and proper. It follows that its image $K(\mu, \nu)$, which we already know is a 1-dimensional submanifold of $\mathcal{T}(S)$, is a line properly embedded in $\mathcal{T}(S)$. $\square$

Following the terminology of [Se1] (motivated by [Ke2]), let the *Kerckhoff line* be the proper 1-dimensional submanifold $K(\mu, \nu) \subset \mathcal{T}(S)$, consisting of all the $\kappa(t\mu, \nu)$ with $t > 0$.

2. NECESSARY CONDITION FOR SMALL BENDING

Let $t \mapsto m_t$, $t \in [0, \varepsilon[$, be a small differentiable curve in $\mathcal{Q}(S)$. If $\beta(m_t) = (\beta^+(t), \beta^-(t)) \in \mathcal{ML}(S)^2$ is its bending measured geodesic lamination, it is shown in [Bo3] that the right derivative $\frac{d}{dt}\beta(m_t)|_{t=0}$ exists, as an element of the tangent space of $\mathcal{ML}(S)^2$ at $\beta(m_0)$. In general, because $\mathcal{ML}(S)$ is not a differentiable manifold, this tangent space consists of geodesic laminations with a transverse structure which is less regular than a transverse measure [Bo2]. However, if we assume in addition that the starting point m_0 of the curve is fuchsian, the tangent space of $\mathcal{ML}(S)^2$ at $\beta(m_0) = (0, 0)$ is just $\mathcal{ML}(S)^2$; see [Bo2].

We can therefore consider the converse problem: Given a fuchsian metric $m_0 \in \mathcal{F}(S)$ and a pair $(\mu, \nu) \in \mathcal{ML}(S)^2$ of measured geodesic laminations, does there exist a small differentiable curve $t \mapsto m_t \in \mathcal{Q}(S)$, $t \in [0, \varepsilon[$, originating from m_0 and such that $\frac{d}{dt}\beta(m_t)|_{t=0} = (\mu, \nu)$? The following result shows that m_0 is completely determined by μ and ν .

Note that, by construction, there is a natural identification between the submanifold $\mathcal{F}(S) \subset \mathcal{Q}(S)$ consisting of all fuchsian metrics and the Teichmüller space $\mathcal{T}(S)$.

Proposition 6. *Let $\mu, \nu \in \mathcal{ML}(S)$ be two measured geodesic laminations, and let $t \mapsto m_t$, $t \in [0, \varepsilon[$ be a differentiable curve in $\mathcal{Q}(S)$, originating from a fuchsian metric m_0 and such that the derivative $\frac{d}{dt}\beta(m_t)|_{t=0}$ of the bending measured lamination is equal to (μ, ν) . Then μ and ν fill up the surface S , and $m_0 \in \mathcal{F}(S) = \mathcal{T}(S)$ is equal to the minimum $\kappa(\mu, \nu)$ of the length function $l_\mu + l_\nu : \mathcal{T}(S) \rightarrow \mathbb{R}$.*

Proof. We consider two other curves in $\mathcal{Q}(S)$.

The first one is the *pure bending* $t \mapsto B_{m_0}^{t\mu}$, obtained by bending the surface S along the measured geodesic lamination $t\mu$ while keeping the metric induced on this pleated surface equal to m_0 . For $t \geq 0$ small enough, $B_{m_0}^{t\mu}$ is a quasifuchsian metric for which the positive side of the boundary $\partial C(B_{m_0}^{t\mu})$ is a pleated surface with induced metric m_0 and with bending measured geodesic lamination $t\mu$. See [EpM, §3] or [Bo1] for the construction of $B_{m_0}^{t\mu}$, and [Mar, §9] to guarantee that it is quasifuchsian for t sufficiently small. In addition, it is proved in [EpM, §3.9] [Bo1] that this curve is differentiable in $\mathcal{Q}(S)$, and in particular admits a tangent vector $b_{m_0}^\mu = \frac{d}{dt}B_{m_0}^{t\mu}|_{t=0} \in T_{m_0}\mathcal{Q}(S)$. This tangent vector $b_{m_0}^\mu$ is the *infinitesimal pure bending* of $m_0 \in \mathcal{F}(S)$ along the measured geodesic lamination μ .

The second curve will use the shear-bend coordinates associated, as in [Bo1], to a maximal geodesic lamination λ containing the support of μ . These coordinates provide a local parametrization of $\mathcal{Q}(S)$ in terms of the geometry of a pleated surface with pleating locus λ . Let m'_t correspond to a pleated surface whose induced metric is equal to the metric $m_t^+ \in \mathcal{T}(S)$ induced on the positive component of the boundary $\partial C(m_t)$ of the convex core, and whose bending data is equal to $t\mu$. By [Mar, §9], m_t is a quasifuchsian metric for t small.

It is proved in [Bo3] that, because the curve $t \mapsto m_t$ is differentiable and because $\frac{d}{dt}\beta^+(m_t)|_{t=0} = \mu$, the right derivative $\dot{m}_0^+ = \frac{d}{dt}m_t^+|_{t=0} \in T_{m_0}\mathcal{T}(S)$ exists and the two curves $t \mapsto m_t$ and $t \mapsto m'_t$ have the same tangent vector at $t = 0$. In particular, by differentiability of the shear-bend coordinates, the tangent vector $\dot{m}_0 = \frac{d}{dt}m_t|_{t=0} = \frac{d}{dt}m'_t|_{t=0} \in T_{m_0}\mathcal{Q}(S)$ is the sum of $b_{m_0}^\mu$ and of $\dot{m}_0^+ = \frac{d}{dt}m_t^+|_{t=0} \in T_{m_0}\mathcal{T}(S) = T_{m_0}\mathcal{F}(S)$.

Similarly, bending S in the negative direction, we can define the infinitesimal pure bending vector $b_{m_0}^{-\nu} = -b_{m_0}^\nu \in T_{m_0}\mathcal{Q}(S)$ of m_0 along $-\nu$. We can also consider the metric $m_t^- \in \mathcal{T}(S)$ induced on the negative side of $\partial C(m_t)$. Then, as above, the vector $\dot{m}_0$ is the sum of $b_{m_0}^{-\nu}$ and of $\dot{m}_0^- = \frac{d}{dt}m_t^-|_{t=0} \in T_{m_0}\mathcal{T}(S) = T_{m_0}\mathcal{F}(S)$.

Finally, we will use the complex structure of $\mathcal{Q}(S)$ coming from the fact that the isometry group of $\mathbb{H}^3$ is $\mathrm{PSL}_2(\mathbb{C})$. Indeed, considering the holonomy of hyperbolic metrics embeds $\mathcal{Q}(S)$ into the space $\mathcal{R}(S)$ of (conjugacy classes) of representations $\pi_1(S) \rightarrow \mathrm{PSL}_2(\mathbb{C})$. This representation space $\mathcal{R}(S)$ is a complex manifold near the image of $\mathcal{Q}(S)$, and admits this image as a complex submanifold (an open subset if S is compact). The shear-bend local coordinates are well behaved with respect to this complex structure; see [Bo1]. In particular, multiplication by $i = \sqrt{-1}$ exchanges shearing and bending. We will use the following two consequences of this. First of all, at a fuchsian metric m_0 , the tangent space $T_{m_0}\mathcal{Q}(S)$ is the direct sum of $T_{m_0}\mathcal{F}(S)$ and of $iT_{m_0}\mathcal{F}(S)$. In addition, the infinitesimal pure bending vector $b_{m_0}^\mu$ belongs to $iT_{m_0}\mathcal{F}(S)$ and is equal to $ie_{m_0}^\mu$, where $e_{m_0}^\mu \in T_{m_0}\mathcal{T}(S) = T_{m_0}\mathcal{F}(S)$ is the infinitesimal earthquake vector along μ .

Applying the decomposition $T_{m_0}\mathcal{Q}(S) = T_{m_0}\mathcal{F}(S) \oplus iT_{m_0}\mathcal{F}(S)$ to the vector $\dot{m}_0 = \dot{m}_0^+ + b_{m_0}^\mu = \dot{m}_0^- + b_{m_0}^{-\nu}$, we conclude that $b_{m_0}^\mu = b_{m_0}^{-\nu}$. Multiplying by $-i$, it

follows that $e_{m_0}^\mu = e_{m_0}^{-\nu}$. As in the proof of Proposition 3, this is equivalent to the property that $m_0 = \kappa(\mu, \nu)$. $\square$

3. REALIZING SMALL BENDING

The goal of this section is to prove a converse to Proposition 6, by constructing in Proposition 11 a small curve of quasifuchsian metrics $t \mapsto m_t \in \mathcal{Q}(S)$, $t \in [0, \varepsilon[$, such that $\beta(m_t) = (t\mu, t\nu)$ for every t .

For the measured geodesic lamination $\mu \in \mathcal{ML}(S)$, let $\mathcal{P}^+(\mu)$ (resp. $\mathcal{P}^-(\mu)$) be the space of quasifuchsian metrics m such that the positive (resp. negative) component of the convex core boundary $\partial C(m)$ has bending measured geodesic lamination $t\mu$, for some $t \in [0, \infty[$.

Recall that $\theta = -3\chi(S) + p$ denotes the dimension of the Teichmüller space $\mathcal{T}(S)$.

Lemma 7. *The space $\mathcal{P}^\pm(\mu)$ is a submanifold-with-boundary of $\mathcal{Q}(S)$, with dimension $\theta + 1$ and with boundary $\mathcal{F}(S)$.*

Proof. We can use the coordinates developed in [Bo1], and associated to a maximal geodesic lamination λ containing the support of μ . These provide an open differentiable embedding $\varphi : \mathcal{Q}(S) \rightarrow \mathcal{T}(S) \times \mathcal{H}_0(\lambda; \mathbb{R}/2\pi\mathbb{Z})$. The first component of $\varphi(m)$ is the hyperbolic metric induced on the unique m -pleated surface f_m with pleating locus λ . The second component is the bending transverse cocycle of f_m , which belongs to the topological group $\mathcal{H}_0(\lambda; \mathbb{R}/2\pi\mathbb{Z}) \cong (\mathbb{R}/2\pi\mathbb{Z})^\theta \oplus \mathbb{Z}/2$ of all $(\mathbb{R}/2\pi\mathbb{Z})$ -valued transverse cocycles for λ that satisfy a certain cusp condition. In general, the bending of the pleated surface f_m is measured by a transverse cocycle and not by a measured geodesic lamination because f_m is not necessarily locally convex.

For notational convenience, it is useful to lift φ to an embedding $\psi : \mathcal{Q}(S) \rightarrow \mathcal{T}(S) \times \mathcal{H}_0(\lambda; \mathbb{R})$, such that ψ sends $\mathcal{F}(S) \cong \mathcal{T}(S)$ to $\mathcal{T}(S) \times \{0\}$ by the identity. Here, $\mathcal{H}_0(\lambda; \mathbb{R}) \cong \mathbb{R}^\theta$ denotes the space of $\mathbb{R}$ -valued transverse cocycle satisfying the cusp condition. Such a ψ exists and is unique because $\mathcal{Q}(S)$ is simply connected.

The vector space $\mathcal{H}_0(\lambda; \mathbb{R})$ contains the transverse measure (also denoted by μ) of the measured geodesic lamination μ , and therefore also contains the two rays $[0, \infty[\mu$ and $] -\infty, 0]\mu$, consisting of all positive (resp. negative) real multiples of μ .

We claim that $\mathcal{P}^+(\mu)$ locally corresponds, under ψ , to the intersection of $\psi(\mathcal{Q}(S))$ with $\mathcal{T}(S) \times [0, \infty[\mu$. Clearly, ψ sends an element of $\mathcal{P}^+(\mu)$ to $\mathcal{T}(S) \times [0, \infty[\mu$. Conversely, it is proved in [KaT] that the map $\eta : \mathcal{Q}(S) \rightarrow \mathcal{T}(S) \times \mathcal{ML}(S)$ which, to a quasifuchsian metric $m \in \mathcal{Q}(S)$, associates the induced metric $m^+ \in \mathcal{T}(S)$ and the bending measured lamination $\beta^+(m)$ of the positive boundary component of the convex core $C(m)$ is a local homeomorphism. In addition, interpreting the ray $[0, \infty[\mu$ as a subset of both $\mathcal{H}_0(\lambda; \mathbb{R})$ and $\mathcal{ML}(S)$, the local inverse for η constructed in [KaT] coincides with ψ on $\mathcal{T}(S) \times [0, \infty[\mu$. It follows that, if $m \in \mathcal{Q}(S)$ is sufficiently close to $m_0 \in \mathcal{P}^+(\mu)$ and if $\psi(m) \in \mathcal{T}(S) \times [0, \infty[\mu$, the bending measured lamination $\beta^+(m)$ is equal to the second component $t\mu$ of $\psi(m)$. Therefore, a metric m near $m_0 \in \mathcal{P}^+(\mu)$ is in $\mathcal{P}^+(\mu)$ if and only if $\psi(m) \in \mathcal{T}(S) \times [0, \infty[\mu$.

A similar property holds for $\mathcal{P}^-(\mu)$ by symmetry. Therefore, under the diffeomorphism ψ , $\mathcal{P}^+(\mu)$, $\mathcal{P}^-(\mu)$ and $\mathcal{F}(S)$ locally correspond to the intersection of $\psi(\mathcal{Q}(S))$ with $\mathcal{T}(S) \times [0, \infty[\mu$, $\mathcal{T}(S) \times] -\infty, 0]\mu$ and $\mathcal{T}(S) \times \{0\}$, respectively. $\square$

Given two measured geodesic laminations $\mu, \nu \in \mathcal{ML}(S)$, we want to consider the intersection of $\mathcal{P}^+(\mu)$ and $\mathcal{P}^-(\nu)$. Note that this intersection is far from being transverse, since these two $(\theta + 1)$ -dimensional submanifolds both contain the θ -dimensional submanifold $\mathcal{F}(S)$ as their boundary. For this reason, we will consider the manifold-with-boundary $\check{\mathcal{Q}}(S)$ obtained by blowing-up $\mathcal{Q}(S)$ along the submanifold $\mathcal{F}(S)$. Namely, $\check{\mathcal{Q}}(S)$ is the union of $\mathcal{Q}(S) - \mathcal{F}(S)$ and of the unit normal bundle $N^1\mathcal{F}(S)$ with the appropriate topology. Recall that the normal bundle $N\mathcal{F}(S) \rightarrow \mathcal{F}(S)$ is intrinsically defined as the bundle whose fiber $N_m\mathcal{F}(S)$ at $m \in \mathcal{F}(S)$ is the quotient $T_m\mathcal{Q}(S)/T_m\mathcal{F}(S)$, and that the fiber $N_m^1\mathcal{F}(S)$ of the unit normal bundle $N^1\mathcal{F}(S) \rightarrow \mathcal{F}(S)$ is the quotient of $N_m\mathcal{F}(S) - \{0\}$ under the multiplicative action of $\mathbb{R}^+$. Considering a tubular neighborhood of $\mathcal{F}(S)$, $\check{\mathcal{Q}}(S)$ is easily endowed with a natural structure of differentiable manifold with boundary $N^1\mathcal{F}(S)$.

Exploiting the complex structure of $\mathcal{Q}(S)$, identify the normal bundle $N\mathcal{F}(S)$ to $iT\mathcal{F}(S)$. The inclusion map $\mathcal{P}^+(\mu) - \mathcal{F}(S) \rightarrow \mathcal{Q}(S) - \mathcal{F}(S)$ uniquely extends to an embedding $\mathcal{P}^+(\mu) \rightarrow \check{\mathcal{Q}}(S)$, which to $m \in \mathcal{F}(S) = \partial\mathcal{P}^+(\mu)$ associates the unit normal vector $\bar{b}_m^\mu \in N^1\mathcal{F}(S)$ which is in the direction of the infinitesimal pure bending vector b_m^μ of m along $+\mu$. We can similarly define an embedding $\mathcal{P}^-(\nu) \rightarrow \check{\mathcal{Q}}(S)$, by associating to $m \in \mathcal{F}(S)$ the unit normal vector $-\bar{b}_m^\nu \in N^1\mathcal{F}(S)$ which is in the direction of the infinitesimal pure bending vector $b_m^{-\nu} = -b_m^\nu$ of m along $-\nu$. Let $\check{\mathcal{P}}^+(\mu)$ and $\check{\mathcal{P}}^-(\nu) \subset \check{\mathcal{Q}}(S)$ be the respective images of these embeddings.

Lemma 8. *The subspaces $\check{\mathcal{P}}^+(\mu)$ and $\check{\mathcal{P}}^-(\nu)$ are submanifolds of $\check{\mathcal{Q}}(S)$, with boundary contained in $\partial\check{\mathcal{Q}}(S)$.*

Proof. The embedding $\psi : \mathcal{Q}(S) \rightarrow T(S) \times \mathcal{H}_0(\lambda; \mathbb{R})$ of the proof of Lemma 7 sends the vector b_m^μ to the vector $(0, \mu)$ in the tangent space of $T(S) \times \mathcal{H}_0(\lambda; \mathbb{R})$. The result then immediately follows from the fact that ψ locally identifies $\mathcal{P}^+(\mu)$, $\mathcal{P}^-(\mu)$ and $T(S)$ to $T(S) \times [0, \infty[\mu$, $T(S) \times]-\infty, 0]\mu$ and $T(S) \times \{0\}$, respectively. This proves Lemma 7. $\square$

Proposition 9. *The boundaries $\partial\check{\mathcal{P}}^+(\mu)$ and $\partial\check{\mathcal{P}}^-(\nu)$ have a non-empty intersection if and only if μ and ν fill up the surface S . If μ and ν fill up S , the intersection of $\partial\check{\mathcal{P}}^+(\mu)$ and $\partial\check{\mathcal{P}}^-(\nu)$ in $\partial\check{\mathcal{Q}}(S)$ is transverse, and is equal to the image of the Kerckhoff line $K(\mu, \nu)$ under the section $m \mapsto \bar{b}_m^\mu = -\bar{b}_m^\nu \in \partial\check{\mathcal{Q}}(S)$.*

Proof. The diffeomorphism $T^1T(S) = T^1\mathcal{F}(S) \rightarrow iT^1\mathcal{F}(S) = N^1\mathcal{F}(S) = \partial\check{\mathcal{Q}}(S)$ defined by multiplication by i sends $\bar{e}_m^\mu$ to $\bar{b}_m^\mu$ and $\bar{e}_m^{-\nu}$ to $\bar{b}_m^{-\nu}$. This translates Proposition 9 to a simple rephrasing of Proposition 5. $\square$

An immediate consequence of Proposition 9 is that the intersection of the two $(\theta + 1)$ -dimensional submanifolds $\check{\mathcal{P}}^+(\mu)$ and $\check{\mathcal{P}}^-(\nu)$ in $\check{\mathcal{Q}}(S)$ is transverse near the boundary $\partial\check{\mathcal{Q}}(S)$. In particular, the intersection $\check{\mathcal{P}}^+(\mu) \cap \check{\mathcal{P}}^-(\nu)$ is a 2-dimensional submanifold of $\check{\mathcal{Q}}(S)$ near $\partial\check{\mathcal{Q}}(S)$, with boundary $\partial\check{\mathcal{P}}^+(\mu) \cap \partial\check{\mathcal{P}}^-(\nu)$ contained in $\partial\check{\mathcal{Q}}(S)$.

By definition, a metric $m \in \mathcal{P}^+(\mu) \cap \mathcal{P}^-(\nu)$ has bending measured lamination $\beta(m) = (t\mu, u\nu)$ for some $t, u \geq 0$. This gives a differentiable map $\pi : \mathcal{P}^+(\mu) \cap \mathcal{P}^-(\nu) \rightarrow \mathbb{R}^2$, defined by $m \mapsto (t, u)$.

Let $\check{\mathbb{R}}^2$ be obtained by blowing up $\mathbb{R}^2$ along $\{0\}$. Because $\pi^{-1}(0) = \mathcal{F}(S)$, the map π lifts to a differentiable map $\tilde{\pi} : \check{\mathcal{P}}^+(\mu) \cap \check{\mathcal{P}}^-(\nu) \rightarrow \check{\mathbb{R}}^2$.

Lemma 10. *The map $\tilde{\pi} : \check{\mathcal{P}}^+(\mu) \cap \check{\mathcal{P}}^-(\nu) \rightarrow \check{\mathbb{R}}^2$ is a local diffeomorphism near $\partial\check{\mathcal{P}}^+(\mu) \cap \partial\check{\mathcal{P}}^-(\nu)$.*

Proof. We will prove that, at any point $p_0 \in \partial\check{\mathcal{P}}^+(\mu) \cap \partial\check{\mathcal{P}}^-(\nu)$, the (linear) tangent map $T_{p_0}\tilde{\pi} : T_{p_0}\check{\mathcal{P}}^+(\mu) \cap T_{p_0}\check{\mathcal{P}}^-(\nu) \rightarrow T_{\tilde{\pi}(p_0)}\check{\mathbb{R}}^2$ is injective.

Let $v \in T_{p_0}\check{\mathcal{P}}^+(\mu) \cap T_{p_0}\check{\mathcal{P}}^-(\nu)$ be such that $T_{p_0}\tilde{\pi}(v) = 0$. Considering the map $\mathcal{P}^+(\mu) \rightarrow [0, \infty[$ which to $m \in \mathcal{P}^+(\mu)$ associates $\beta^+(m)/\mu \in [0, \infty[$ and the induced map $\check{\mathcal{P}}^+(\mu) \rightarrow [0, \infty[$, we see that v must necessarily be in the tangent space of the boundary $\partial\check{\mathcal{P}}^+(\mu)$. Symmetrically, it must be in $T_{p_0}\partial\check{\mathcal{P}}^-(\nu)$. Therefore, v is tangent to the intersection $\partial\check{\mathcal{P}}^+(\mu) \cap \partial\check{\mathcal{P}}^-(\nu)$.

Let us analyze the restriction of $\tilde{\pi}$ to the boundary $\partial\check{\mathcal{P}}^+(\mu) \cap \partial\check{\mathcal{P}}^-(\nu)$. By Proposition 9, $\partial\check{\mathcal{P}}^+(\mu) \cap \partial\check{\mathcal{P}}^-(\nu)$ is equal to the image of the Kerckhoff line $K(\mu, \nu)$ under the section $m \mapsto \bar{b}_m^\mu = -\bar{b}_m^\nu \in \partial\check{\mathcal{Q}}(S)$. Recall that an element of the Kerckhoff line $K(\mu, \nu)$ is of the form $m = \kappa(t\mu, \nu)$ for some $t > 0$, which is equivalent to the property that $b_m^{t\mu} = -b_m^\nu$. We will also need a coordinate chart for $\check{\mathbb{R}}^2$ near $\tilde{\pi}(p_0)$. Noting that the image of π is contained in the quadrant $[0, \infty[^2$, we can use for this the chart $\varphi :]0, \infty[\times]0, \infty[\rightarrow \check{\mathbb{R}}^2$ defined on the interior by $(x, y) \mapsto (xy, y)$.

We claim that, if $m = \kappa(t\mu, \nu) \in K(\mu, \nu)$, then $\varphi^{-1} \circ \tilde{\pi}(\bar{b}_m^\mu)$ is just equal to $(t, 0)$. To see this, choose, in the 2-dimensional manifold $\check{\mathcal{P}}^+(\mu) \cap \check{\mathcal{P}}^-(\nu)$, a small curve $s \mapsto \check{m}_s$, $s \in [0, \varepsilon[$, such that $\check{m}_0 = \bar{b}_m^\mu \in \partial\check{\mathcal{P}}^+(\mu) \cap \partial\check{\mathcal{P}}^-(\nu)$ and such that $\frac{d}{ds}\check{m}_s|_{s=0}$ is not tangent to the boundary. The curve $s \mapsto \check{m}_s$ projects to a differentiable curve $t \mapsto m_t \in \mathcal{Q}(S)$ with $m_0 = m$. By definition of $\mathcal{P}^+(\mu)$ and $\mathcal{P}^-(\nu)$, the bending measured lamination $\beta(m_s)$ is of the form $(t(s)\mu, u(s)\nu)$ for two differentiable functions $t(s)$, and $u(s)$ with $t(0) = u(0) = 0$. Since $\frac{d}{ds}\check{m}_s|_{s=0}$ points away from the boundary, the curve $s \mapsto m_s$ is not tangent to $\mathcal{F}(S)$ at $s = 0$, and it follows that at least one of the derivatives $t'(0)$, $u'(0)$ is non-trivial. If we apply Proposition 6, we conclude that $m = \kappa(t'(0)\mu, u'(0)\nu)$. Since $m = \kappa(t\mu, \nu)$, it follows that $t'(0)/u'(0) = t$ by Lemma 4. In particular, $t(s)/u(s)$ tends to t as s tends to 0. Therefore, $\varphi^{-1} \circ \tilde{\pi}(\bar{b}_m^\mu)$, which is the limit of $\varphi^{-1} \circ \tilde{\pi}(\check{m}_s) = \varphi^{-1} \circ \pi(m_s) = \varphi^{-1} \circ (t(s), u(s)) = (t(s)/u(s), u(s))$ as $s > 0$ tends to 0, is equal to $(t, 0)$.

This computation shows that the restriction of $\tilde{\pi}$ to the boundary $\partial\check{\mathcal{P}}^+(\mu) \cap \partial\check{\mathcal{P}}^-(\nu)$ is a diffeomorphism onto its image. In particular, if $v \in T_{p_0}\partial\check{\mathcal{P}}^+(\mu) \cap T_{p_0}\partial\check{\mathcal{P}}^-(\nu)$ is such that $T_{p_0}\tilde{\pi}(v) = 0$, then necessarily $v = 0$.

This concludes the proof that the tangent map $T_{p_0}\tilde{\pi} : T_{p_0}\check{\mathcal{P}}^+(\mu) \cap T_{p_0}\check{\mathcal{P}}^-(\nu) \rightarrow T_{\tilde{\pi}(p_0)}\check{\mathbb{R}}^2$ is injective. Since $\tilde{\pi}$ sends the boundary of the 2-dimensional manifold $\check{\mathcal{P}}^+(\mu) \cap \check{\mathcal{P}}^-(\nu)$ to the boundary of the 2-dimensional manifold $\check{\mathbb{R}}^2$, this proves that $\tilde{\pi} : \check{\mathcal{P}}^+(\mu) \cap \check{\mathcal{P}}^-(\nu) \rightarrow \check{\mathbb{R}}^2$ is a local diffeomorphism near $p_0 \in \partial\check{\mathcal{P}}^+(\mu) \cap \partial\check{\mathcal{P}}^-(\nu)$. $\square$

This immediately gives the following converse to Proposition 6.

Proposition 11. *Let $\mu, \nu \in \mathcal{ML}(S)$ be two measured geodesic laminations which fill up the surface S , and let m_0 be the minimum $\kappa(\mu, \nu)$ of the length function $l_\mu + l_\nu$. Then there is a small differentiable curve $t \mapsto m_t \in \mathcal{Q}(S)$, $t \in [0, \varepsilon[$, beginning at m_0 and such that the bending measured lamination $\beta(m_t)$ is equal to $(t\mu, t\nu)$ for every t .*

Proof. Consider the curve $t \mapsto (t, t)$, $t \in [0, \varepsilon[$, in $\mathbb{R}^2$. By Lemma 10, for ε small enough, there is a curve $t \mapsto \check{m}_t \in \check{\mathcal{P}}^+(\mu) \cap \check{\mathcal{P}}^-(\nu)$ such that $t \mapsto \tilde{\pi}(\check{m}_t)$ coincides

with the lift of $t \mapsto (t, t)$ to $\check{\mathbb{R}}^2$. By definition of the map π , this just means that the projection $m_t \in \mathcal{Q}(S)$ of $\tilde{m}_t \in \check{\mathcal{Q}}(S)$ is such that $\beta(m_t) = (t\mu, t\nu)$. $\square$

4. PARAMETRIZING QUASI-FUCHSIAN GROUPS BY THEIR SMALL BENDING

Recall that $\mathcal{FML}(S)$ denotes the open subset of $\mathcal{ML}(S)^2$ consisting of those pairs (μ, ν) such that μ and ν fill up the surface S .

Let $\check{\mathcal{FML}}(S)$ be obtained by blowing up $\mathcal{FML}(S) \cup \{(0, 0)\}$ along $\{(0, 0)\}$. Namely, $\check{\mathcal{FML}}(S)$ is formally obtained from $\mathcal{FML}(S)$ by extending each ray $]0, \infty[(\mu, \nu)$ to a semi-open ray $[0, \infty[(\mu, \nu)$, with the obvious topology. Note that the boundary $\partial\check{\mathcal{FML}}(S)$ is just the quotient space of $\mathcal{FML}(S)$ under the multiplicative action of $\mathbb{R}^+$.

For every $(\mu, \nu) \in \mathcal{FML}(S)$, Proposition 11 provides a maximal ray $R_{\mu\nu} = [0, \varepsilon_{\mu\nu}[(\mu, \nu)$ in $\mathcal{FML}(S) \cup \{(0, 0)\}$ and a differentiable map $\Phi_{\mu\nu} : R_{\mu\nu} \rightarrow \mathcal{Q}(S)$ such that $\Phi_{\mu\nu}(\mu', \nu')$ has bending measured lamination (μ', ν') for every $(\mu', \nu') \in R_{\mu\nu}$ and such that $\check{\mathcal{P}}^+(\mu)$ and $\check{\mathcal{P}}^-(\nu)$ meet transversely along $\beta(R_{\mu\nu})$. Here, the statement that $R_{\mu\nu}$ is maximal means that $\varepsilon_{\mu\nu} \in]0, \infty]$ is maximal for this property.

Note that $R_{\mu\nu}$ and $\varphi_{\mu\nu}$ depend only on the orbit of (μ, ν) under the action of $\mathbb{R}^+$, namely on the corresponding point of $\partial\check{\mathcal{FML}}(S)$. Let $\check{R}_{\mu\nu}$ be the lift of $R_{\mu\nu}$ in $\check{\mathcal{FML}}(S)$, and lift $\Phi_{\mu\nu}$ to $\check{\Phi}_{\mu\nu} : \check{R}_{\mu\nu} \rightarrow \check{\mathcal{Q}}(S)$. In particular, $\check{\Phi}_{\mu\nu}$ sends the initial point of $\check{R}_{\mu\nu}$ to the bending vector $\check{b}_m^\mu = \check{b}_m^{-\nu} \in N^1\mathcal{F}(S) = \partial\check{\mathcal{Q}}(S)$ with $m = \kappa(\mu, \nu)$.

Let $\check{U} \subset \check{\mathcal{FML}}(S)$ denote the union of all the $\check{R}_{\mu\nu}$, and let $\check{\Phi} : \check{U} \rightarrow \mathcal{FML}(S)$ restrict to $\check{\Phi}_{\mu\nu}$ on each $\check{R}_{\mu\nu}$. Note that the property that $\Phi_{\mu\nu}(\mu', \nu')$ has bending measured lamination (μ', ν') implies that the $\check{R}_{\mu\nu}$ are pairwise disjoint, so that $\check{\Phi}$ is well-defined. We want to show that $\check{\Phi}$ is continuous.

Lemma 12. *As the measured geodesic lamination μ tends to μ_0 for the topology of $\mathcal{ML}(S)$, the submanifold $\mathcal{P}^\pm(\mu)$ tends to $\mathcal{P}^\pm(\mu_0)$ for the topology of C^∞ convergence on compact subsets.*

Proof. We will use the tools developed in [Bo1].

Let $\mu_n \in \mathcal{ML}(S)$, $n \in \mathbb{N}$, be a sequence converging to μ_0 . Let λ_n be a maximal geodesic lamination containing the support of μ_n . Passing to a subsequence if necessary, we can assume that, for the Hausdorff topology, the geodesic lamination λ_n converges to a geodesic lamination λ_0 , which is necessarily maximal and contains the support of μ_0 .

The shear-bend coordinates associated to λ_n provide an open biholomorphic embedding $\Phi_n : \mathcal{Q}(S) \rightarrow \mathcal{H}_0(\lambda_n; \mathbb{C}/2\pi i\mathbb{Z})$. Here $\mathcal{H}_0(\lambda_n; \mathbb{C}/2\pi i\mathbb{Z})$ is the topological group of $\mathbb{C}/2\pi i\mathbb{Z}$ -valued transverse cocycles for the maximal geodesic lamination λ_n which satisfy the cusp condition, and is isomorphic to $(\mathbb{C}/2\pi i\mathbb{Z})^\theta \oplus \mathbb{Z}/2$. For a metric $m \in \mathcal{Q}(S)$, the real part of $\Phi_n(m) \in \mathcal{H}_0(\lambda_n; \mathbb{C}/2\pi i\mathbb{Z}) = \mathcal{H}_0(\lambda_n; \mathbb{R}) \oplus i\mathcal{H}_0(\lambda_n; \mathbb{R}/2\pi\mathbb{Z})$ measures the induced metric of the unique m -pleated surface with bending locus λ_n , and the imaginary part measures its bending. In particular, $\mathcal{P}^\pm(\mu_n)$ locally corresponds to $\Phi_n^{-1}(\mathcal{H}_0(\lambda_n; \mathbb{R}) \oplus i\mathbb{R}\mu_n)$, or more precisely to a branch of the immersion which is the composition of the projection $\mathcal{H}_0(\lambda_n; \mathbb{R}) \oplus i\mathbb{R}\mu_n \rightarrow \mathcal{H}_0(\lambda_n; \mathbb{C}/2\pi i\mathbb{Z})$ and of Φ_n^{-1} .

To compare the various $\mathcal{H}_0(\lambda_n; \mathbb{C}/2\pi i\mathbb{Z})$ pick a train track τ carrying λ_0 . Since λ_n converges to λ_0 for the Hausdorff topology, τ also carries the λ_n for n large enough. Then there is a well-defined isomorphism $\Psi_n : \mathcal{H}_0(\lambda_n; \mathbb{C}/2\pi i\mathbb{Z}) \rightarrow$

$\mathcal{W}(\tau; \mathbb{C}/2\pi i\mathbb{Z})$, where $\mathcal{W}(\tau; \mathbb{C}/2\pi i\mathbb{Z})$ is the group of $\mathbb{C}/2\pi i\mathbb{Z}$ -valued edge weights for τ satisfying the switch and cusp relations.

Because λ_n converges to λ_0 for the Hausdorff topology, it follows from the explicit construction of [Bo1, §5, §8] that $\Phi_n^{-1} \circ \Psi_n^{-1}$ converges to $\Phi_0^{-1} \circ \Psi_0^{-1}$, uniformly on compact subsets of the image of $\Psi_0 \circ \Phi_0$. Because these maps are holomorphic, the convergence is actually C^∞ . Since μ_n converges to μ_0 for the topology of $\mathcal{ML}(S)$, the edge weight system $\Psi_n(\mu_n)$ converges to $\Psi_0(\mu_0)$ in $\mathcal{W}_0(\tau, \mathbb{R})$. It follows that $\mathcal{P}^\pm(\mu_n)$, which locally corresponds to $\Phi_n^{-1} \circ \Psi_n^{-1}(\mathcal{W}_0(\tau; \mathbb{R}) \oplus i\mathbb{R}\Psi_n(\mu_n))$, converges to $\mathcal{P}^\pm(\mu_0)$, which locally corresponds to $\Phi_0^{-1} \circ \Psi_0^{-1}(\mathcal{W}_0(\tau; \mathbb{R}) \oplus i\mathbb{R}\Psi_0(\mu_0))$, in the topology of C^∞ -convergence on compact subsets. $\square$

Note that we actually proved real analytic convergence in Lemma 12. However, we will only need C^2 convergence.

Theorem 13. *The subset $\tilde{U}$ is an open neighborhood of $\partial\tilde{\mathcal{F}}\mathcal{ML}(S)$ in $\tilde{\mathcal{F}}\mathcal{ML}(S)$, and $\tilde{\Phi}$ is a homeomorphism from $\tilde{U}$ to an open neighborhood $\tilde{V}$ of $\partial\tilde{\mathcal{Q}}(S)$ in $\tilde{\mathcal{Q}}(S)$.*

Proof. The restriction $\tilde{\Phi}_{\mu\nu}$ of $\tilde{\Phi}$ to $\tilde{R}_{\mu\nu}$ was constructed by considering the transverse intersection of the submanifolds $\tilde{\mathcal{P}}^+(\mu)$ and $\tilde{\mathcal{P}}^-(\nu)$ near the boundary of $\tilde{\mathcal{Q}}(S)$. By Lemma 12, $\tilde{\mathcal{P}}^+(\mu)$ and $\tilde{\mathcal{P}}^-(\nu)$ depend continuously on (μ, ν) for the topology of C^1 convergence. (Note that one needs the C^2 continuity of $\mathcal{P}^+(\mu)$ and $\mathcal{P}^-(\nu)$ to guarantee the C^1 continuity of $\tilde{\mathcal{P}}^+(\mu)$ and $\tilde{\mathcal{P}}^-(\nu)$ near the boundary.) It follows that the length $\varepsilon_{\mu\nu}$ of $R_{\mu\nu} = [0, \varepsilon_{\mu\nu}[$ is a lower semi-continuous function of (μ, ν) , and that $\tilde{\Phi}_{\mu\nu}$ depends continuously on (μ, ν) . This proves that the union $\tilde{U}$ of the $\tilde{R}_{\mu\nu}$ is open in $\partial\tilde{\mathcal{F}}\mathcal{ML}(S)$, and that $\tilde{\Phi}$ is continuous.

Let $p : \tilde{\mathcal{Q}}(S) \rightarrow \mathcal{Q}(S)$ be the natural projection and, as usual, let $\beta : \mathcal{Q}(S) \rightarrow \mathcal{ML}(S)^2$ be the bending map. By construction, $\beta \circ p \circ \tilde{\Phi}$ is the identity map on $\tilde{U} - \partial\tilde{\mathcal{F}}\mathcal{ML}(S) \subset \mathcal{ML}(S)^2$. It follows that $\tilde{\Phi}$ is injective on $\tilde{U} - \partial\tilde{\mathcal{F}}\mathcal{ML}(S)$, and therefore on all of $\tilde{U}$ by Proposition 6.

The two spaces $\tilde{U} \subset \tilde{\mathcal{F}}\mathcal{ML}(S)$ and $\tilde{\mathcal{Q}}(S)$ are topological manifolds-with-boundary of the same dimension 2θ . The map $\tilde{\Phi} : \tilde{U} \rightarrow \tilde{\mathcal{F}}\mathcal{ML}(S)$ is continuous and injective, and sends boundary points to boundary points. By the Theorem of Invariance of the Domain, it follows that its image $\tilde{V} = \tilde{\Phi}(\tilde{U})$ is open in $\tilde{\mathcal{Q}}(S)$, and that $\tilde{\Phi}$ restricts to a homeomorphism $\tilde{U} \rightarrow \tilde{V}$. $\square$

Theorem 14. *There exists an open neighborhood V of the fuchsian submanifold $\mathcal{F}(S)$ in $\mathcal{Q}(S)$ such that the bending map $\beta : \mathcal{Q}(S) \rightarrow \mathcal{ML}(S)^2$ induces a homeomorphism between $V - \mathcal{F}(S)$ and its image.*

Proof. For the canonical identifications between $\tilde{\mathcal{Q}}(S) - \partial\tilde{\mathcal{Q}}(S)$ and $\mathcal{Q}(S) - \mathcal{F}(S)$, and between $\tilde{\mathcal{F}}\mathcal{ML}(S) - \partial\tilde{\mathcal{F}}\mathcal{ML}(S)$ and $\mathcal{F}\mathcal{ML}(S)$, the inverse of the restriction of $\tilde{\Phi}$ to $\tilde{U} - \partial\tilde{\mathcal{Q}}(S)$ coincides with β . Therefore, the only part which requires some checking is that the image V of $\tilde{V} = \tilde{\Phi}(\tilde{U})$ under the canonical projection $p : \tilde{\mathcal{Q}}(S) \rightarrow \mathcal{Q}(S)$ is open in $\mathcal{Q}(S)$. However, this immediately follows from the fact that the preimages of points under p are all compact, which implies that p is an open map. (Note that this is false for the projection $\tilde{\mathcal{F}}\mathcal{ML}(S) \rightarrow \mathcal{F}\mathcal{ML}(S) \cup \{0\}$.) $\square$

Theorem 14 is just Theorem 1 stated in the introduction. Theorem 2 immediately follows from the definition of $\tilde{U}$ and from the fact that $\tilde{U}$ is open (Theorem 13).

We conclude with a few remarks.

A recent result of Caroline Series [Se2] shows that we can restrict the neighborhood V of Theorem 14 so that $\beta^{-1}(\beta(V)) = V$.

When μ and ν are multicurves, namely when their supports consist of finitely many closed geodesics, it follows from [HoK] that the submanifolds $\check{\mathcal{P}}^+(\mu)$ and $\check{\mathcal{P}}^-(\nu)$ are everywhere transverse, by a doubling argument as in [BoO]. Consequently, the open subset V of Theorem 14 can be chosen so that the image $U = \beta(V) \subset \mathcal{ML}(S)^2$ contains all rays of the form $[0, \infty[(\mu, \nu)$ where μ and ν are multicurves.

As indicated in the introduction, it is conjectured that we can take V equal to the whole space $\mathcal{Q}(S)$. See [BoO] for a characterization of the image of $\mathcal{Q}(S)$ under β .

REFERENCES

- [Bo1] Francis Bonahon, *Shearing hyperbolic surfaces, bending pleated surfaces and Thurston's symplectic form*, Ann. Fac. Sci. Toulouse Math. (6) **5** (1996), pp. 233–297.
- [Bo2] ———, *Geodesic laminations with transverse Hölder distributions*, Ann. Sci. École Norm. Sup. (4) **30** (1997), pp. 205–240.
- [Bo3] ———, *The boundary geometry of 3-dimensional hyperbolic convex cores*, J. Differential Geometry **50** (1998), pp. 1–23.
- [BoO] Francis Bonahon, Jean-Pierre Otal, *Laminations mesurées de plissage des variétés hyperboliques de dimension 3*, preprint, 2001.
- [EpM] David B. A. Epstein, Albert Marden, *Convex hulls in hyperbolic spaces, a theorem of Sullivan, and measured pleated surfaces*, in: *Analytical and geometric aspects of hyperbolic space* (D.B.A. Epstein ed.), L.M.S. Lecture Note Series vol. **111**, 1986, Cambridge University Press, pp. 113–253.
- [HoK] Craig D. Hodgson, Steven P. Kerckhoff, *Rigidity of hyperbolic cone-manifolds and hyperbolic Dehn surgery*, J. Differential Geom. **48** (1998), pp. 1–59.
- [KaT] Yoshinobu Kamishima, Ser P. Tan, *Deformations spaces of geometric structures*, in: *Aspects of low-dimensional manifolds* (Y. Matsumo, S. Morita eds.), Advanced Studies in Pure Math. **20**, 1992, Kinokuniya Company Ltd., Tokyo, pp. 263–299.
- [Ke1] Steven P. Kerckhoff, *The Nielsen realization problem*, Ann. of Math. **117** (1983), pp. 235–265.
- [Ke2] ———, *Lines of minima in Teichmüller space*, Duke Math. J. **65** (1992), pp. 187–213.
- [Mar] Albert Marden, *The geometry of finitely generated Kleinian groups*, Ann. of Math. **99** (1974), pp. 383–462.
- [Mas] Bernard Maskit, *Kleinian groups*, Grundlehren der Math. Wiss. **287**, Springer-Verlag, 1988.
- [Se1] Caroline M. Series, *On Kerckhoff minima and pleating loci for quasi-Fuchsian groups*, Geom. Dedicata **88** (2001), pp. 211–237.
- [Se2] ———, *Limits of quasifuchsian groups with small bending*, preprint, 2002.
- [Su] Dennis P. Sullivan, *Quasiconformal homeomorphisms and dynamics II: Structural stability implies hyperbolicity for Kleinian groups*, Acta Math. **155** (1985), pp. 243–260.
- [Th1] William P. Thurston, *The topology and geometry of 3-manifolds*, Lecture notes, Princeton University, 1976–79.
- [Th2] ———, *Earthquakes in two-dimensional hyperbolic geometry*, in: *Low-dimensional Topology and Kleinian groups* (D.B.A. Epstein ed.), L.M.S. Lecture Notes Series vol. **112**, 1986, Cambridge University press, pp. 91–112.
- [Wo1] Scott A. Wolpert, *On the symplectic geometry of deformations of a hyperbolic surface*, Ann. of Math. **117** (1983), pp. 207–234.
- [Wo2] ———, *Geodesic length functions and the Nielsen problem*, J. Differential Geom. **25** (1987), pp. 275–296.

F. BONAHON, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTHERN CALIFORNIA, LOS ANGELES, CA 90089-1113, U.S.A.

E-mail address: `fbonahon@math.usc.edu`

URL: `http://math.usc.edu/~fbonahon`